



### PRODUCT DESCRIPTION

- RIVERINA TURKEY STARTER CRUMBLES are suitable to be fed as a complete feed for growing turkeys from day old through to approximately seven (7) weeks of age. Turkey poults should have unlimited access (feed Ad lib.) to RIVERINA TURKEY STARTER CRUMBLES.
- This feed is the first stage in a two stage-feeding programme for growing turkeys. It is produced in a fine crumble form so that feed wastage is minimised, digestibility is improved and so that a consistent intake of nutrients essential for growth is achieved. This feed is formulated in accordance with current research into turkey nutrition and is mixed from a variety of quality raw materials.

### COMPOSITION

- Base grain as depicted seasonally, and therefore proportions may vary.
- Wheat, maize, sorghum, bran and pollard, a combination of protein meals (cottonseed meal, soyabean meal, sunflower meal, blood meal, meat meal), vegetable/animal oil, limestone, sodium bicarbonate, salt, choline chloride, amino acids, Avatec™, Riverina Vitamin and Minerals.

### DIRECTIONS FOR USE

- The first three to five weeks are of vital importance as rearing turkey poults. Upon arrival of the new day old poults, it may be necessary to dip the beaks of some of them in feed and water to encourage the consumption of both of these.
- This feed is a complete feed and no supplementary feeding is necessary.
- During this period approximate feed intakes should be:

| Age of Birds | Approximate Feed Consumption (per head per day) |
|--------------|---|
| Week One     | 10-20 grams                                     |
| Week Two     | 25-35 grams (toms)<br>20-25 grams (hens)        |
| Week Three   | 60-65 grams (toms)<br>55-60 grams (hens)        |
| Week Four    | 85-90 grams (toms)<br>65-70 grams (hens)        |
| Week Five    | 100-105 grams (toms)<br>85-90 grams (hens)      |
| Week Six     | 120-130 grams (toms)<br>105-115 grams (hens)    |
| Week Seven   | 155-165 grams (toms)<br>125-135 grams (hens)    |

- NOTE: The above feed intakes are only indicative. Feed intakes will vary with the season, variety of bird, lighting, shed conditions, disease status and other environmental conditions.
- For the first two days of feeding it is best to spread some feed on trays or newspaper to enable all birds' access.
- It is important to change the feed regularly so it remains fresh and palatable.
- It is very important that birds have clean, cool, fresh water available at all times.

### TECHNICAL RELEVANCE

- Protein meals providing natural sources of essential amino acids. Plus amino acids to balance essential nutrient requirements.
- Contains essential vitamins and minerals.
- Turkey poults are susceptible to diseases such as coccidiosis. Medication has been included in this feed to aid in the prevention of coccidiosis.

### VETERINARY CHEMICALS:

- Avatec™ as Lasalocid sodium added at the rate of 95 mg/kg.

### MEDICINAL CLAIM:

This feed contains the following veterinary chemicals-

- Avatec™ (lasalocid sodium) is added to this feed for the prevention of coccidiosis caused by Eimeria tenella, E. acervulina, E. maxima and E. brunetti.

### NUTRITIONAL ANALYSIS

|            |       |      |           |
|------------|-------|------|-----------|
| PROTEIN    | %     | :    | 28.00     |
| CALCIUM    | %     | :    | 0.8-1.2   |
| AV PHOS    | %     | MIN: | 0.40      |
| ENERGYA-ME | MJ/kg | :    | 11.65     |
| SALT       | %     | MIN  | 0.15-0.40 |

### SPECIAL FEATURES

#### IN CRUMBLE FORM

- Eliminates separation that can occur with mash/meal type feeds. Improved digestibility. Consistent nutrient intake.

#### QUALITY PRODUCT

- Manufactured to strict quality control standards.

#### PRECISE FORMULATION AND MIXING

- Formulated in accordance with current research into poultry nutrition and mixed from a variety of quality raw materials.

### WITHOLDING PERIOD

Meat: Nil. Eggs: Do not use in birds laying eggs for human consumption.

### WARNING

After withdrawal of medicated feed, monitor flock for coccidiosis outbreaks and treat with Avatec™ Coccidiostat solution. Coccidiosis control programs may need to be designed for a particular flock, property or situation and veterinary advice may be necessary.

*Please note Government legislation prohibits the inclusion of medication used for blackhead. Should the incidence of Blackhead occur, please contact your local veterinarian for advice on medication suitable for blackhead control.*

### STORAGE

Cool, dry conditions, away from vermin.

### EXPIRY AFTER MANUFACTURE

3 months from the date of manufacture.

### PACK SIZE

20kg woven polypropylene bags.

Or

Bulk loads: Contact your local Riverina Sales Representative for details.

Avatec™ is a product of OzBioPharm P/L.



This feed is scientifically designed for the intended species of animal and should not be fed to any other animal. It may contain medication and ingredients that may prove harmful if fed to other species.  
Ver: 3.0

Issue Date: Oct 2007

